

## The One-Round Voronoi Game\*

Otfried Cheong,<sup>1</sup> Sarel Har-Peled,<sup>2</sup> Nathan Linial,<sup>3</sup> and Jiří Matoušek<sup>4</sup>

<sup>1</sup>Department of Mathematics and Computer Science, Eindhoven University of Technology,  
P.O. Box 513, 5600 MB Eindhoven, The Netherlands  
o.cheong@tue.nl

<sup>2</sup>Department of Computer Science, University of Illinois,  
1304 W. Springfield Ave., Urbana, IL 61801, USA  
sariel@cs.uiuc.edu

<sup>3</sup>Institute of Computer Science, Hebrew University,  
Jerusalem 91904, Israel  
nati@cs.huji.ac.il

<sup>4</sup>Department of Applied Mathematics and  
Institute of Theoretical Computer Science (ITI), Charles University,  
Malostranské nám. 25, 118 00 Praha 1, Czech Republic  
matousek@kam.mff.cuni.cz

**Abstract.** In the one-round Voronoi game, the first player chooses an  $n$ -point set  $\mathcal{W}$  in a square  $Q$ , and then the second player places another  $n$ -point set  $\mathcal{B}$  into  $Q$ . The payoff for the second player is the fraction of the area of  $Q$  occupied by the regions of the points of  $\mathcal{B}$  in the Voronoi diagram of  $\mathcal{W} \cup \mathcal{B}$ . We give a (randomized) strategy for the second player that always guarantees him a payoff of at least  $\frac{1}{2} + \alpha$ , for a constant  $\alpha > 0$  and every large enough  $n$ . This contrasts with the one-dimensional situation, with  $Q = [0, 1]$ , where the first player can always win more than  $\frac{1}{2}$ .

### 1. Introduction

*Competitive facility location* studies the placement of sites by competing market players. Overviews of different models are the surveys by Tobin et al. [9], Eiselt and Laporte [3], and Eiselt et al. [4].

---

\* Part of this research was done during a workshop supported by ITI (Project LN00A056 of the Ministry of Education of the Czech Republic). N.L. was supported in part by a grant from the Israel Science Foundation.

The *Voronoi game* is a simple geometric model for competitive facility location, where a site  $s$  “owns” the part of the playing arena that is closer to  $s$  than to any other site. We consider a two-player version with a square arena  $Q$ . The two players, White and Black, place points into  $Q$ . As in chess, White plays first. The goal of both players is to capture as much of the area of  $Q$  as possible, where the region captured by White is

$$R(\mathcal{W}, \mathcal{B}) = \{x \in Q : \text{dist}(x, \mathcal{W}) < \text{dist}(x, \mathcal{B})\}$$

and the region captured by Black is  $R(\mathcal{B}, \mathcal{W})$ . Here  $\mathcal{W}$  is the set of points of White,  $\mathcal{B}$  is the set of points of Black, and  $\text{dist}(\cdot, \cdot)$  is the Euclidean distance. In other words, if we construct the Voronoi diagram of  $\mathcal{W} \cup \mathcal{B}$ , then each player captures the Voronoi regions (restricted to  $Q$ ) of his point set and is rewarded proportionate to the measure of his captured set. The *payoff* for White is

$$\frac{\text{vol}(R(\mathcal{W}, \mathcal{B}))}{\text{vol}(Q)}$$

and the payoff for Black is

$$\frac{\text{vol}(R(\mathcal{B}, \mathcal{W}))}{\text{vol}(Q)},$$

where  $\text{vol}(\cdot)$  is the Lebesgue measure. (Of course, we can rescale the arena  $Q$  so that  $\text{vol}(Q) = 1$ , but in the subsequent considerations a different scaling seems more intuitive.)

Ahn et al. [1] studied a one-dimensional Voronoi game, where the arena  $Q$  is a line segment, and the game takes  $n$  rounds. In each round, White and Black place one point each. Ahn et al. showed that Black then has a winning strategy that guarantees a payoff of  $\frac{1}{2} + \varepsilon$ , with  $\varepsilon > 0$ , but that White can force  $\varepsilon$  to be as small as he wishes. On the other hand, if only a single round is played, where White first places  $n$  points, followed by Black placing  $n$  points, then White has a winning strategy. In fact, if  $Q = [0, 2n]$  and White plays on the odd integer points  $\{1, 3, 5, \dots, 2n - 1\}$ , then Black’s payoff is less than  $\frac{1}{2}$ .

In this paper we show that in the two-dimensional case Black, rather than White, has a winning strategy: for each set  $\mathcal{W}$  of  $n$  points, there is a set  $\mathcal{B}$  of  $n$  black points such that Black’s payoff is at least  $\frac{1}{2} + \alpha$ , for an absolute constant  $\alpha > 0$  and  $n$  large enough.

From now on, let  $Q$  be the square  $[0, \sqrt{n}]^2$ , of area  $n$ , so that the average area per white point is 1. To win the game, Black needs to find  $n$  points such that their average area is at least  $\frac{1}{2} + \alpha$ . We first show that it is very easy to find *one* such point—in fact, a *random* point in  $Q \setminus \mathcal{W}$  has this property. Since this is the key idea of our proof, we first present it in a modified setting where the arena  $Q$  has the topology of a torus, eliminating boundary effects. We then proceed to prove this result for the square with its standard topology, showing how to handle the square boundary, and proceed to prove the result for  $n$  black points. Finally, we generalize the result to higher dimensions.

After a preliminary version of the present paper appeared in the *Proceedings of the 18th ACM Symposium on Computational Geometry* (2002), further progress on the one-round Voronoi game was made by Fekete and Meijer [7]. In particular, for rectangular

arenas  $Q$  they showed that White has a winning strategy if and only if  $n = 1$ , or  $n = 2$  and  $\rho \leq \sqrt{3}/2$ , or  $n \geq 3$  and  $\rho \leq \sqrt{2}/n$ , where  $\rho \leq 1$  is the ratio of the sides of  $Q$ , and Black has a winning strategy in all other cases (a winning strategy need not imply winning by a fixed margin).

## 2. The Torus Case

To present the (simple) main idea of our proofs in a setting free of technical complications due to effects near the boundary of  $Q$ , we assume in this section that the square  $Q$  has the topology of a torus. To be precise, we identify the left and right edges of  $Q$ , as well as the top and bottom edges, and we alter the Euclidean metric in  $Q$  accordingly.

**Proposition 1.** *There exist constants  $\beta > 0$  and  $n_0$  such that for every  $n$ -point set  $\mathcal{W}$  in the square arena  $Q$  with torus topology,  $n \geq n_0$ , there is a point  $x \in Q \setminus \mathcal{W}$  with*

$$\text{vol}(R(x, \mathcal{W})) \geq \frac{1}{2} + \beta.$$

*In fact,  $x$  can be selected uniformly at random:*

$$\mathbf{E}[\text{vol}(R(x, \mathcal{W}))] \geq \frac{1}{2} + \beta,$$

*where  $\mathbf{E}[\cdot]$  denotes expectation with respect to uniform random selection of  $x \in Q$ .*

*Proof.* If there is a point  $p \in Q$  such that  $\text{dist}(p, \mathcal{W}) > \sqrt{n}/4$ , then the proposition holds: with probability bounded below by a positive constant, the point  $x$  will grab an  $\Omega(n)$  area, which is definitely larger than (say) 1 for sufficiently large  $n$ . In the following we can therefore assume that no such point  $p$  exists. In this case each region in the Voronoi diagram of  $\mathcal{W}$  is topologically a disk, and the dual graph (where two points  $p, q \in \mathcal{W}$  are connected by an edge if their Voronoi regions have a common edge) has no multiple edges. Any graph without multiple edges drawn on a torus has average degree at most 6 (see for instance Proposition 4.4.4 of [8]), a fact that will be useful later.

Let  $I_A$  denote the characteristic function of a set  $A$ . We have

$$\begin{aligned} \mathbf{E}[\text{vol}(R(x, \mathcal{W}))] &= \frac{1}{\text{vol}(Q)} \int_Q \int_Q I_{R(x, \mathcal{W})}(y) \, dy \, dx \\ &= \frac{1}{n} \int_Q \text{vol}(\{x \in Q : y \in R(x, \mathcal{W})\}) \, dy \end{aligned}$$

by Fubini's theorem.

A point  $y \in Q$  lies in  $R(x, \mathcal{W})$  if and only if  $\text{dist}(y, x) < r = \text{dist}(y, \mathcal{W})$ , and so

$$\{x \in Q : y \in R(x, \mathcal{W})\} = \{x \in Q : \text{dist}(x, y) < r\}.$$

Since  $r \leq \sqrt{n}/4$ , this is a disk of radius  $r$  centered at  $y$  (possibly wrapping around the edges of  $Q$ ).

Our integral thus becomes  $(\pi/n) \int_Q \text{dist}(y, \mathcal{W})^2 dy$ , a quantity that we denote by  $F_0(\mathcal{W})$ . We split it into integrals over  $\mathcal{W}$ 's Voronoi cells:

$$F_0(\mathcal{W}) = \frac{\pi}{n} \sum_{w \in \mathcal{W}} \int_{\text{cell}_{\mathcal{W}}(w)} \text{dist}(y, w)^2 dy,$$

where  $\text{cell}_{\mathcal{W}}(w)$  is the region of  $w$  in the Voronoi diagram of  $\mathcal{W}$  in  $Q$ .

Among all convex bodies  $C \subset \mathbf{R}^2$  of area  $a$ , the integral  $\int_C \text{dist}(y, w)^2 dy$  is minimized by the disk  $C_0$  of area  $a$  centered at  $w$  (somewhat informally, moving a piece of  $C$  closer to  $w$  decreases the integral, and such a move is possible for any  $C$  but that disk). The value of that integral over  $C_0$  is

$$\int_{C_0} \text{dist}(y, w)^2 dy = \int_0^{\sqrt{a/\pi}} r^2 \cdot 2\pi r dr = \frac{a^2}{2\pi}.$$

Moreover, we will need the following intuitively obvious lemma:

**Lemma 2.** *For every  $k \geq 3$  there exists a constant  $\varepsilon_k > 0$  such that if  $C$  is a convex  $k$ -gon of area  $a$ , then  $\int_C \text{dist}(y, w)^2 dy \geq (1 + \varepsilon_k) \int_{C_0} \text{dist}(y, w)^2 dy = (1 + \varepsilon_k)(a^2/2\pi)$ .*

A formal proof for arbitrary dimension will be given as Lemma 7 in Section 5.

We set  $a_w := \text{vol}(\text{cell}_{\mathcal{W}}(w))$ . Then

$$\begin{aligned} F_0(\mathcal{W}) &= \frac{\pi}{n} \sum_{w \in \mathcal{W}} \int_{\text{cell}_{\mathcal{W}}(w)} \text{dist}(y, w)^2 dy \\ &\geq \frac{1}{2n} \sum_{w \in \mathcal{W}} a_w^2 \geq \frac{1}{2n} \frac{\left(\sum_{w \in \mathcal{W}} a_w\right)^2}{n} \geq \frac{1}{2} \end{aligned}$$

by Cauchy–Schwarz.

We see that for a random point  $x$ , the expected region size is at least  $\frac{1}{2}$ , but we want  $\frac{1}{2} + \beta$ . By Lemma 2, if  $\text{cell}_{\mathcal{W}}(w)$  has at most  $k$  sides, then  $\int_{\text{cell}_{\mathcal{W}}(w)} \text{dist}(y, w)^2 dy \geq (1 + \varepsilon_k) \cdot (a_w^2/2\pi)$ . Let  $\mathcal{W}_f \subseteq \mathcal{W}$  consist of the points whose regions in the Voronoi diagram of  $\mathcal{W}$  have fewer than 12 sides. Since, as was mentioned earlier, the average number of sides of a cell in the Voronoi diagram of  $\mathcal{W}$  is at most 6, we have  $|\mathcal{W}_f| \geq \frac{1}{2}n$ .

So we win the factor  $1 + \varepsilon_{11}$  in at least half of the regions and lose nothing in the other regions. The only problem is that the regions of  $\mathcal{W}_f$  could together occupy only a tiny fraction of the area of  $Q$  and then this win would not reach the threshold  $\beta > 0$  that we seek.

Let us assume that they occupy, say, less than  $\frac{1}{4}$  of the total area. Then the average area of the remaining regions (of  $\mathcal{W} \setminus \mathcal{W}_f$ ) is at least  $\frac{3}{2}$  (at most  $\frac{1}{2}n$  regions take up area at least  $\frac{3}{4}n$ ). The Cauchy–Schwarz inequality used in the calculation above then becomes strict and we win a constant factor in the regions of  $\mathcal{W} \setminus \mathcal{W}_f$ . Namely,

$$F_0(\mathcal{W}) \geq \frac{1}{2n} \sum_{w \in \mathcal{W} \setminus \mathcal{W}_f} a_w^2$$

$$\begin{aligned}
 &\geq \frac{1}{2n} \frac{(\sum_{w \in \mathcal{W} \setminus \mathcal{W}_f} a_w)^2}{|\mathcal{W} \setminus \mathcal{W}_f|} \\
 &\geq \frac{1}{2n} \cdot \frac{3}{2} \cdot \sum_{w \in \mathcal{W} \setminus \mathcal{W}_f} a_w \geq \frac{1}{2n} \cdot \frac{3}{2} \cdot \frac{3}{4} n \geq \frac{9}{16}. \quad \square
 \end{aligned}$$

### 3. The Proof with Boundary Effects

The torus arena conveniently removed the need to consider the boundary effects. We now prove the same result for the square with boundary:

**Proposition 3.** *There exist constants  $\beta > 0$  and  $n_0$  such that for every  $n$ -point set  $\mathcal{W} \subset Q$ ,  $n \geq n_0$ , we have*

$$\mathbf{E}[\text{vol}(R(x, \mathcal{W}))] \geq \frac{1}{2} + \beta.$$

*Proof.* As in the proof of Proposition 1, we can rewrite the expected area as

$$\begin{aligned}
 F(\mathcal{W}) &= \frac{1}{n} \int_Q \text{vol}(\{x \in Q : y \in R(x, \mathcal{W})\}) \, dy \\
 &= \frac{1}{n} \int_Q \text{vol}(B(y, \text{dist}(y, \mathcal{W})) \cap Q) \, dy \\
 &= \frac{1}{n} \sum_{w \in \mathcal{W}} \int_{\text{cell}_{\mathcal{W}}(w)} \text{vol}(B(y, \text{dist}(y, w)) \cap Q) \, dy,
 \end{aligned}$$

where  $B(x, r)$  is the disk of radius  $r$  centered at  $x$ . We want to bound  $F(\mathcal{W})$  from below by  $\frac{1}{2} + \beta$ .


We choose a large constant  $D$  (the requirements on  $D$  will become apparent later). We call a region  $\text{cell}_{\mathcal{W}}(w)$  *long* if it has diameter at least  $D$  and *short* otherwise, and we denote by  $\mathcal{W}_\ell$  and  $\mathcal{W}_s$  the subsets of  $\mathcal{W}$  corresponding to the long and short regions, respectively. Let again  $a_w := \text{vol}(\text{cell}_{\mathcal{W}}(w))$ .

First we consider the long regions. We note that for any  $w, y \in Q$ ,

$$\text{vol}(B(y, \text{dist}(y, w)) \cap Q) \geq \frac{1}{2} \cdot \text{dist}(y, w)^2 \tag{1}$$

(the extreme case is  $w$  and  $y$  in opposite corners of  $Q$ ).

Now let  $w \in \mathcal{W}_\ell$  and write  $C = \text{cell}_{\mathcal{W}}(w)$ . We claim that at least  $\frac{1}{16}$  of the area of  $C$  lies at distance at least  $\frac{1}{4}D$  from  $w$ ; in other words,  $\text{vol}(C \setminus B(w, \frac{1}{4}D)) \geq \frac{1}{16}a_w$  (the constant can be improved). Let  $p, q$  be a diametrical pair of points of  $C$ , and place two copies  $C_p, C_q$  of  $\frac{1}{4}C$  inside  $C$  so that they share a common tangent to  $C$  at  $p$  and  $q$ , respectively, where  $\frac{1}{4}C$  is the shape resulting from shrinking  $C$  by a factor of 4; see Fig. 1. Clearly, the distance between  $C_p$  and  $C_q$  is  $D/2$ , and consequently, either  $C_p$  or  $C_q$  do not intersect  $B = B(w, \frac{1}{4}D)$ . Thus, the area of  $C$  not covered by  $B$  is at least  $\text{vol}(C_p) = \text{vol}(C_q) = \text{vol}(C)/16$ .



**Fig. 1.** At least  $\frac{1}{16}$  of the area of  $C$  is not covered by  $B(w, \frac{1}{4}D)$ .

It follows that

$$\int_{\text{cell}_{\mathcal{W}}(w)} \text{vol}(B(y, \text{dist}(y, w)) \cap Q) \, dy \geq \frac{1}{2} \cdot \frac{D^2}{16} \cdot \frac{1}{16} a_w > \frac{D^2}{1000} a_w$$

for every  $w \in \mathcal{W}_\ell$ , and so the contribution of the long regions to  $F(\mathcal{W})$  is at least  $(D^2/1000n)A_\ell$ , where  $A_\ell = \sum_{w \in \mathcal{W}_\ell} a_w$ .

Next, we consider the short regions (of diameter at most  $D$ ), and among those only the *inner* ones, whose distance to the boundary of  $Q$  is at least  $D$ . Let  $\mathcal{W}_{\text{si}}$  be the corresponding subset of  $\mathcal{W}$ . We have  $A_{\text{si}} = \sum_{w \in \mathcal{W}_{\text{si}}} a_w \geq n - 8D\sqrt{n} - A_\ell$ . For the short inner regions, the disk  $B(y, \text{dist}(y, w))$  lies completely inside  $Q$  and so their contribution to  $F(\mathcal{W})$  behaves as in the proof of Proposition 1; it equals

$$\frac{\pi}{n} \sum_{w \in \mathcal{W}_{\text{si}}} \int_{\text{cell}_{\mathcal{W}}(w)} \text{dist}(y, w)^2 \, dy.$$

As we saw above, this quantity is bounded below by

$$\frac{1}{2n} \sum_{w \in \mathcal{W}_{\text{si}}} a_w^2 \geq \frac{1}{2n} \frac{A_{\text{si}}^2}{|\mathcal{W}_{\text{si}}|}.$$

Now we distinguish several cases depending on the orders of magnitude of  $A_\ell$  and  $|\mathcal{W}_{\text{si}}|$ .

- (1) Suppose that  $A_\ell \geq n/2D$ . Then the contribution of  $A_\ell$  alone suffices:  $F(\mathcal{W}) \geq (D^2/1000n)A_\ell \geq D/2000 > \frac{1}{2} + \beta$  for  $D$  large enough.
- (2) Now let  $A_\ell < n/2D$ . For large  $n$  this implies  $A_{\text{si}} \geq (1 - 1/D)n$ . Now two cases are distinguished according to  $|\mathcal{W}_{\text{si}}|$ .
  - (a) For  $|\mathcal{W}_{\text{si}}| \leq (1 - 4/D)n$ , we obtain

$$F(\mathcal{W}) \geq \frac{1}{2n} \frac{A_{\text{si}}^2}{|\mathcal{W}_{\text{si}}|} \geq \frac{(1 - 1/D)^2}{2(1 - 4/D)} \geq \frac{1}{2} + \frac{1}{D},$$

which is the desired bound.

- (b) It remains to deal with the case  $A_{\text{si}} \geq (1 - 1/D)n$  and  $|\mathcal{W}_{\text{si}}| \geq (1 - 4/D)n$ . If  $D$  is very large, we are essentially in the situation analyzed in the proof of Proposition 1. We can repeat the argument from that proof: We consider the subset  $\mathcal{W}_{\text{sif}} \subseteq \mathcal{W}_{\text{si}}$  consisting of points whose regions have at most 12 sides, we get that  $\mathcal{W}_{\text{sif}}$  has almost  $\frac{1}{2}n$  points, and we win a fixed constant factor for each of the regions of  $\mathcal{W}_{\text{sif}}$ , etc. The inequalities that were exactly true in the proof of Proposition 1 may now hold only up to multiplicative factors that tend to 1 as  $D \rightarrow \infty$ , but everything goes through and we get  $F(\mathcal{W}) \geq \frac{1}{2} + \beta$  in this case as well. We omit the tedious detailed calculations.  $\square$

#### 4. The Main Result

A key ingredient in the proof of our main theorem is the following lemma, showing that if Black throws in  $\delta n$  points at random, instead of one as in Proposition 3, then his expected area gain still exceeds  $\frac{1}{2}\delta n$  at least by a fixed fraction, provided that  $\delta > 0$  is sufficiently small.

**Lemma 4.** *For every sufficiently large constant  $D$ , there exist constants  $\beta_1 > 0$ ,  $\delta > 0$ , and  $n_0$  such that for every  $n$ -point set  $\mathcal{W} \subset Q$ ,  $n \geq n_0$ , if  $\mathcal{B} \subset Q$  is obtained by  $\delta n$  independent random draws from the uniform distribution on  $Q$ , then*

$$\mathbf{E}[\text{vol}(R(\mathcal{B}, \mathcal{W}))] \geq (\frac{1}{2} + \beta_1)\delta n.$$

*If the total area  $A_\ell$  of the long regions (of diameter at least  $D$ ) exceeds  $n/2D$ , then*

$$\mathbf{E}[\text{vol}(R(\mathcal{B}, \mathcal{W}))] \geq 2\delta n.$$

*Proof.* This is very similar to the proof of Proposition 3. Intuitively, for small  $\delta$ , the  $\delta n$  independent random points are likely to interact very little and their expected area gain is likely to be nearly  $(\delta - O(\delta^2))n$  times the expected area gain of a single point.

This time we have

$$\mathbf{E}[\text{vol}(R(\mathcal{B}, \mathcal{W}))] = \int_Q \text{Prob}[y \in R(\mathcal{B}, \mathcal{W})] \, dy.$$

Here  $P(y) = \text{Prob}[y \in R(\mathcal{B}, \mathcal{W})]$  is the probability with respect to the random choice of the set  $\mathcal{B}$ . Namely,


$$\begin{aligned} P(y) &= \text{Prob}[\mathcal{B} \cap B(y, \text{dist}(y, \mathcal{W})) \neq \emptyset] \\ &= 1 - (\text{Prob}[x \notin B(y, \text{dist}(y, \mathcal{W}))])^{\delta n} \\ &= 1 - \left(1 - \frac{1}{n} \cdot \text{vol}(B(y, \text{dist}(y, \mathcal{W})) \cap Q)\right)^{\delta n}. \end{aligned}$$

We write  $\rho(y) = (1/n) \cdot \text{vol}(B(y, \text{dist}(y, \mathcal{W})) \cap Q)$ . If  $y$  lies in a short region of the Voronoi diagram of  $\mathcal{W}$ , then  $\rho(y) \leq C_D/n$  with  $C_D$  depending only on  $D$ , and  $\delta C_D$  can be made as small as desired by choosing  $\delta$  sufficiently small. Then we obtain

$$P(y) = 1 - (1 - \rho(y))^{\delta n} \geq \delta n \rho(y) + O((\delta n \rho(y))^2) \geq \delta n \rho(y) \cdot (1 - \gamma)$$

with  $\gamma$  a small constant. Thus, the contribution of a short Voronoi region to  $\mathbf{E}[\text{vol}(R(\mathcal{B}, \mathcal{W}))]$  is at least  $(1 - \gamma)\delta n$  times the contribution of that region to the expected area gained by a single random point as in Proposition 3. All the calculations involving short regions can be done in exactly the same way. It remains to show that if the total area  $A_\ell$  of the long regions is at least  $n/2D$ , then these regions contribute at least  $2\delta n$  to  $\mathbf{E}[\text{vol}(R(\mathcal{B}, \mathcal{W}))]$ .

In the proof of Proposition 3, equation (1), we have shown  $\rho(y) \geq (1/2n) \cdot \text{dist}(y, w)^2$  for  $y \in \text{cell}_{\mathcal{W}}(w)$ . We also know that  $\text{dist}(y, w) \geq \frac{1}{4}D$  for  $y$  in at least  $\frac{1}{16}$  of the area of



**Fig. 2.** Two points of Black can take over almost a complete cell of White.

each long region. For these  $y$ , we have  $P(y) \geq 1 - e^{-\rho(y)\delta n} \geq 1 - e^{-D^2\delta/200} \geq D^2\delta/400$  (assuming  $\delta < D^{-2}$ ). The integral over all the long regions together is then at least  $\frac{1}{16}A_\ell \geq n/32D$  times this quantity, and therefore larger than  $2\delta n$  with ample room to spare.  $\square$

We can now prove our main theorem.

**Theorem 5.** *There exist constants  $\alpha > 0$  and  $n_0$  such that for every  $n \geq n_0$ , Black can always win at least  $\frac{1}{2} + \alpha$  in the Voronoi game. That is, for every  $n$ -point set  $\mathcal{W} \subset Q$  there exists an  $n$ -point set  $\mathcal{B} \subset Q \setminus \mathcal{W}$  with  $\text{vol}(R(\mathcal{B}, \mathcal{W})) \geq (\frac{1}{2} + \alpha) \text{vol}(Q)$ .*

*Proof.* Let  $w \in \mathcal{W}$ . A takeover of  $w$ 's region means that Black places two of his points very close to  $w$  with  $w$  as the center of symmetry; see Fig. 2. In this way he captures almost all of  $\text{cell}_{\mathcal{W}}(w)$ . This suggests the following strategy for Black: a takeover of the  $\frac{1}{2}n$  largest White regions guarantees Black a payoff arbitrarily close to  $\frac{1}{2}n$ . This does not prove the theorem, in general, but it fails to do so only if almost all of White's regions have almost the same area. Thus, if more than  $\varepsilon n$  White regions have area below  $1 - \varepsilon$ , for some constant  $\varepsilon > 0$ , then the takeover strategy implies the theorem. It therefore suffices to describe a strategy<sup>1</sup> for Black when all but  $\varepsilon n$  of White's regions have area at least  $1 - \varepsilon$ .

First Black chooses a set  $\mathcal{B}_0$  of  $\delta n$  points as in Lemma 4; that is, with  $\text{vol}(R(\mathcal{B}_0, \mathcal{W})) \geq (1 + \beta_1)\delta n$  and even with  $\text{vol}(R(\mathcal{B}_0, \mathcal{W})) \geq 2\delta n$  if  $A_\ell \geq n/2D$ .

If  $A_\ell \geq n/2D$ , then White now has  $n$  regions of total area  $A_{\mathcal{W}} \leq (1 - 2\delta)n$  and Black still has  $(1 - \delta)n$  points to place. He takes over the  $\frac{1}{2}(1 - \delta)n$  largest among the current regions of White. In this way Black has captured at least an area arbitrarily close to

$$n - A_{\mathcal{W}} + \frac{1}{2}(1 - \delta)n \cdot \frac{A_{\mathcal{W}}}{n} = n - \frac{1}{2}(1 + \delta)A_{\mathcal{W}} > \frac{1}{2}(1 + \delta)n.$$

Next, we suppose that  $A_\ell < n/2D$ . We consider a point  $w \in \mathcal{W}_s$  defining a short region and call  $w$  contaminated if Black has captured some point of  $\text{cell}_{\mathcal{W}}(w)$  by the set  $\mathcal{B}_0$ . A short region can be contaminated only by a point  $b \in \mathcal{B}_0$  if  $\text{dist}(b, w) \leq 2D$ . Therefore, the total area of contaminated short regions is  $O(D^2\delta n) < n/3$ , say, and so

<sup>1</sup> A similar trick would also simplify the proof of Proposition 3 if we did not want to prove the claim about a random point but only the existence of a point capturing at least  $\frac{1}{2} + \beta$ .



regions of total area at least  $n/2$  remain uncontaminated. Now we use the assumption that all but  $\varepsilon n$  of White's regions have area at least  $1 - \varepsilon$ . Black can now take over the  $\frac{1}{2}(1 - \delta)n$  largest uncontaminated regions. This implies that the number of uncontaminated regions of size  $\geq 1 - \varepsilon$  is at least  $n/2 - \varepsilon n$ . Thus, Black can now occupy at least  $\min(n/2 - \varepsilon n, (1 - \delta)n/2) \geq \frac{1}{2}(1 - \delta)n - \varepsilon n$  cells, to gain total area at least

$$\begin{aligned} & \left(\frac{1}{2} + \beta_1\right)\delta n + \left(\frac{1}{2}(1 - \delta)n - \varepsilon n\right)(1 - \varepsilon) \\ &= \left(\frac{1}{2} + \beta_1\right)\delta n + \frac{1}{2}(1 - \delta - 2\varepsilon)(1 - \varepsilon)n. \end{aligned}$$

If  $\varepsilon$  is very small compared with  $\delta$  and  $\beta_1$ , then the expression above is at least  $(\frac{1}{2} + \alpha)n$  with  $\alpha$  close to  $\beta_1\delta$ . This concludes the proof of the theorem.  $\square$

### 5. The Higher-Dimensional Case

The proof of Proposition 1 (and therefore of Lemma 4) exploited the fact that the Voronoi diagram is a planar graph, and therefore at least half of all Voronoi cells have at most 11 edges. In higher dimensions, though, the average number of facets of a Voronoi cell cannot be bounded by any constant, and so we must argue differently in order to show that the Voronoi cells cannot be all arbitrarily similar to a ball.

**Definition 6.** A convex body  $C$  is  $(1 + \mu)$ -spherical with center  $p$ , for  $\mu > 0$ , if there exists a radius  $r > 0$  such that  $B(p, r) \subseteq C \subseteq B(p, r(1 + \mu))$ .

**Lemma 7.** If a convex body  $C$  in  $\mathbb{R}^d$  is not  $(1 + \mu)$ -spherical with center  $p$ , for some  $p \in \mathbb{R}^d$  and  $\mu > 0$ , then there exists a constant  $\beta > 0$  depending only on  $\mu$  and  $d$  such that

$$\int_C c_d \cdot \text{dist}(y, p)^d \, dy \geq (1 + \beta)L,$$

with

$$L = \int_{\Delta} c_d \cdot \text{dist}(y, p)^d \, dy = \frac{\text{vol}(C)^2}{2}.$$

Here  $\Delta$  is a ball of the same volume as  $C$  centered at  $p$ , and  $c_d$  is the volume of the unit ball in  $\mathbb{R}^d$ .

*Proof.* Let  $\Delta = B(p, R)$ , where  $R = (\text{vol}(C)/c_d)^{1/d}$ . Then

$$\begin{aligned} L &= \int_{\Delta} c_d \cdot \text{dist}(y, p)^d \, dy = \int_0^R (c_d r^d) \cdot (c_d \, dr^{d-1}) \, dr \\ &= \frac{c_d^2}{2} R^{2d} = \frac{\text{vol}(C)^2}{2}. \end{aligned}$$

As for the other claim, let  $r'$  (resp.  $R'$ ) be the largest (resp. smallest) radius so that  $B(p, r') \subseteq C \subseteq B(p, R')$ . Since  $C$  is not  $(1 + \mu)$ -spherical with center  $p$ , it follows

that there exists a positive constant  $\beta_1$  such that  $(1 + \beta_1)R \leq R'$ . In particular, this implies that there exists a constant  $\beta_2$ , such that  $\text{vol}(K) \geq \beta_2 \text{vol}(C)$ , where  $K = C \setminus B(p, R(1 + \beta_1/4))$ . Namely,

$$\begin{aligned} \int_C c_d \cdot \text{dist}(y, p)^d dy &\geq \int_\Delta c_d \cdot \text{dist}(y, p)^d dy + \int_K c_d \left( \left( R \left( 1 + \frac{\beta_1}{4} \right) \right)^d - R^d \right) dy \\ &\geq (1 + \beta) \int_\Delta c_d \cdot \text{dist}(y, p)^d dy = (1 + \beta)L, \end{aligned}$$

where  $\beta > 0$  is an appropriate constant that depends only on  $d$  and  $\mu$ .  $\square$

**Lemma 8.** *Let  $Q$  be a hypercube in  $\mathbb{R}^d$ , and let  $P$  be a set of points in  $Q$ . Let  $V(P)$  denote the decomposition of  $Q$  into convex cells by the Voronoi diagram of  $P$  restricted to  $Q$ . Then there exists a constant  $\mu > 0$ , which depends only on  $d$ , such that the total volume of the cells that are  $(1 + \mu)$ -spherical with respect to their defining site in  $P$  is bounded by  $\text{vol}(Q)/2$ .*

*Proof.* Consider a cell  $C_p$  of  $V(P)$  that is  $(1 + \mu)$ -spherical with center  $p$ . Let  $B(p, r)$  be the largest ball with center  $p$  that is contained inside  $C_p$ . Let  $N(p)$  be the set of points of  $P$  whose Voronoi cells have a common boundary with  $C_p$ .


We claim that all the points of  $N(p)$  lie inside a “thin” spherical shell centered at  $p$ , and, furthermore, the points of  $N(p)$  are dense inside this spherical shell. This implies that their Voronoi cells must be long and skinny, and as such they cannot be  $\mu$ -spherical, for  $\mu$  small enough.

Indeed, observe that the distance of any point of  $N(p)$  to  $p$  is at least  $2r$  and at most  $2r(1 + \mu)$ . Furthermore, any angular cone of angular angle  $4\sqrt{\mu}$  emanating from  $p$  must include a point of  $N(p)$ . Indeed, consider such a cone  $Z$  with a ray  $\rho$  as its rotational axis and angular radius  $4\sqrt{\mu}$ , where  $\rho$  emanates from  $p$ . Let  $s$  denote the intersection of  $\rho$  with the spherical shell  $B(p, (1 + \mu)r) \setminus B(p, r)$ . Since one endpoint of  $s$  is outside  $C_p$ , and the other is inside  $C_p$ , it follows that there must be a point  $q \in P$ , so that the bisector of  $p$  and  $q$  intersects  $s$ . See Fig. 3.

We claim that  $q$  is inside  $Z$ . Indeed, the angle  $\alpha$  between  $pq$  and  $\rho$  is maximized when the bisector between  $p$  and  $q$  is tangent, at a point  $v$ , to  $B(p, r)$ , and it passes through the far endpoint  $u$  of  $s$ . Clearly,  $\angle pvu = \pi/2$ ,  $\text{dist}(p, v) = r$ ,  $\text{dist}(p, u) = (1 + \mu)r$ , and  $\text{dist}(u, v) = r\sqrt{(1 + \mu)^2 - 1} = r\sqrt{2\mu + \mu^2} \leq r\sqrt{3\mu}$ , for  $\mu$  small enough. Now,  $\sin(\alpha) = \text{dist}(u, v)/\text{dist}(p, u) \leq \sqrt{3\mu}/(1 + \mu) \leq 2\sqrt{\mu}$ , for  $\mu$  small enough. Finally,  $\alpha \leq 2 \sin(\alpha) \leq 4\sqrt{\mu}$ , for  $\sin(\alpha)$  small enough. Implying that  $q \in Z$ . See Fig. 4.

This implies that  $N(p)$  is dense. Indeed, consider a point  $t \in N(p)$ . Its nearest point in  $N(p)$  is at distance at most  $2r(1 + \mu) \cdot 2 \cdot 4\sqrt{\mu} = O(r\sqrt{\mu})$ . On the other hand, the Voronoi cell  $C_t$  of  $t$  has a point on its boundary at distance  $\geq r$  from  $t$  (as it shares a boundary point  $u'$  with  $C_p$ ,  $\text{dist}(u', p) \geq r$ , and  $\text{dist}(u', t) = \text{dist}(u', p)$ ). (This also implies that  $C_p$  is not adjacent to the boundary of  $Q$ , as this would imply that some of the points of  $N(p)$  lie outside  $Q$ .)

That is,  $C_q$  is not  $(1 + \gamma)$ -spherical, where  $\gamma = \Omega(r/r\sqrt{\mu} - 1) = \Omega(1/\sqrt{\mu})$ . By making  $\mu$  small enough, we can ensure that  $C_q$  is not  $(1 + \mu)$ -spherical with center  $q$ .




**Fig. 3.** (a) If the Voronoi cell  $C_p$  is  $(1 + \mu)$ -spherical with center  $p$ , then the neighboring sites must be in a spherical shell around  $p$ . (b) The sites of  $N(p)$  are densely spread in this spherical shell, since there must be a site inside the intersection  $\Delta$  between the spherical shell and any cone of angular radius  $4\sqrt{\mu}$ .

We have shown that every  $(1 + \mu)$ -spherical cell in  $V(P)$  is surrounded by cells that are not  $(1 + \mu)$ -spherical. We will charge the volume of such a  $(1 + \mu)$ -spherical cell to its surrounding cells as follows. For a point  $p \in P$  whose Voronoi cell is  $(1 + \mu)$ -spherical with center  $p$ , let  $r_p$  be the radius of the largest ball contained inside  $C_p$  centered at  $p$ , and let  $U_p = B(p, 1.8r_p)$  be the *region of influence* of  $p$ . Clearly,  $U_p \cap P = \{p\}$  and  $\text{vol}(U_p) \geq (1.8/(1 + \mu))^d \text{vol}(C_p) \geq 2 \text{vol}(C_p)$ , for  $\mu$  sufficiently small. By picking  $\mu$  small enough, we can also guarantee that the regions of influence of the  $(1 + \mu)$ -spherical cells of  $V(P)$  are disjoint. We charge the volume of a  $(1 + \mu)$ -spherical cell to its region of influence, establishing the claim.  $\square$

**Theorem 9.** *There exist constants  $\alpha > 0$  and  $n_0$  depending only on the dimension  $d$ , such that for every  $n \geq n_0$ , Black can always win at least  $\frac{1}{2} + \alpha$  in the Voronoi game played on arena  $Q$ , the  $d$ -dimensional hypercube of volume  $n$ . That is, for every  $n$ -point set  $\mathcal{W} \subset Q$  there exists an  $n$ -point set  $\mathcal{B} \subset Q \setminus \mathcal{W}$  with  $\text{vol}(R(\mathcal{B}, \mathcal{W})) \geq (\frac{1}{2} + \alpha) \text{vol}(Q)$ .*

*Proof.* The argumentation follows the two-dimensional case closely, so we sketch it only. First, we consider  $Q$  to have the topology of a torus, by identifying any two extreme faces, in the same dimension, of  $Q$  with each other. Using Lemma 8, we know that at least half of  $Q$  is covered by cells of the Voronoi diagram of  $\mathcal{W}$  which are not “round.” As such, following the argumentation of Proposition 1 and using Lemma 7, we get that for a random point  $x$ , we have  $\mathbf{E}[\text{vol}(R(x, \mathcal{W}))] \geq \frac{1}{2} + \beta$ , where  $\beta$  is an appropriate constant.



**Fig. 4.** Illustration of the proof of Lemma 8.

As in the two-dimensional case, our next step is to remove the assumption about the torus topology. The proof of Proposition 3 can be followed closely. We omit the straightforward details. (Note that as the dimension increases, the boundary effect is more pronounced. However, taking  $n_0$  to be large enough, this has no impact on the proof, except for making the constants somewhat worse.)

Finally, the proofs of Lemma 4 and Theorem 5 are “dimension independent” and apply verbatim in the higher-dimensional case.  $\square$

## 6. Conclusions and Open Problems

We considered the Voronoi game on a square or hypercube board  $Q$ , played in a single round: White starts by placing  $n$  points  $\mathcal{W}$  in  $Q$ , then Black places another  $n$  points  $\mathcal{B}$  disjoint from  $\mathcal{W}$ , and finally the winner is determined.

Our considerations appear to generalize without much change to sufficiently “fat” convex arenas in the plane. On the other hand, when the arena degenerates to a line segment, we have reached the one-dimensional case where White, not Black, has a winning strategy [1]. Fekete and Meijer [7] have analyzed the behavior of the game as function of the aspect ratio of the rectangle, as we mentioned in the Introduction. Their analysis is strongly dependent on the arena being a rectangle, and it would be interesting to see if similar results can be obtained for more general arenas.

What happens when the number of points played by White and Black are not identical? Specifically, let  $\lambda$  be a real number between 0 and 2. Consider the game where White plays  $n$  points and Black plays  $\lambda n$  points. Let  $f(\lambda, n)$  be the payoff to Black in this Voronoi game. It is not hard to show that  $f(0, n) = 0$  and that  $\lim_{n \rightarrow \infty} f(2, n) = 1$ . We know that  $f(\lambda, n) > (\frac{1}{2} + \varepsilon)\lambda$  for some positive  $\varepsilon$  and  $n$  large enough, as long as  $\lambda$  is bounded away from 0 and 2. It would be interesting to get a better idea of the behavior of  $f$ . Does  $\lim_{n \rightarrow \infty} f(\lambda, n)$  exist for all  $\lambda$ ?

We have shown that for any set of  $n$  white points, there is a black point that grabs a “large” Voronoi cell. It would be interesting to find configurations of the white points for which no black point can do too well. Obvious candidates are grid arrangements of the white points, such as the square grid or hexagonal grid.

In fact, if we ask for a configuration of the white points that minimizes the payoff of a *random* black point, it is known that the hexagonal grid is optimal if  $n$  is large enough. This follows from a result on the two-dimensional quantizer problem. In the quantizer problem we want to quantize two-dimensional input values from a continuous domain (a ball  $B \subset \mathbf{R}^2$ , say) using  $\log n$  bits. This is done by choosing a discrete quantizer set  $P$  of  $n$  points in  $B$ , and replacing the input value  $x \in B$  by the closest point from  $P$ . Assuming uniform distribution of the input values, the *mean squared error* of a quantizer  $P$  is

$$\frac{1}{\text{vol}(B)} \sum_{p \in P} \int_{\text{cell}(p)} \text{dist}(x, p)^2 dx,$$

where  $\text{cell}(p)$  is the Voronoi cell of  $p$  in the Voronoi diagram of  $P$  (see [2]). Fejes

Tóth [5] (see also [6]) showed that if  $n$  is sufficiently large, then the error is minimized by choosing  $P$  to be the hexagonal grid.

In the proof of Proposition 1, we showed that the expected payoff of a random black point is

$$\frac{\pi}{n} \sum_{p \in \mathcal{W}} \int_{\text{cell}(w)} \text{dist}(x, w)^2 dx,$$

with the slight twist that here we assume torus topology. Assuming  $n$  is so large that we can ignore the difference in topology, this is proportional to the quantization error of  $\mathcal{W}$ , and so Fejes Tóth's result implies that the optimal choice of  $\mathcal{W}$  is the hexagonal grid. An interesting open question is whether the hexagonal grid is also optimal if we consider the *maximum* possible area that a black point can grab.

The original version of the Voronoi game [1] is played in more than one round: White and Black alternate placing points on the board  $Q$ . The value of this game and the optimal strategies are still unknown for dimensions higher than one. If the arena  $Q$  is centrally symmetric, but the symmetry has no fixed point in  $Q$ , then Black can respond to each move of White with a point placed in the symmetric location. This guarantees a payoff of  $\frac{1}{2}$ . Many obvious questions remain open: Can Black actually win the game for large  $n$ ? What happens with asymmetric boards?

We note that our analysis is somewhat sloppy in relation to the constants. Of specific interest is the margin in which the second player wins over the first player. Currently, it seems that our techniques are too limited to yield any bound remotely close to the truth. In particular, a more promising venue for estimating those constants is either by simulation, or by investigating special configurations (that is, points placed on a lattice).

Finally, we believe that the strategy of the Black player in the one-round Voronoi game suggested by our proofs can be derandomized (yielding a polynomial-time algorithm) using a sliding grid argument. We have not elaborated this in detail, though.

## Acknowledgment

We thank three anonymous referees for careful reading and for many useful comments.

## References

1. H. Ahn, S. Cheng, O. Cheong, M. Golin, and R. van Oostrum. Competitive facility location along a highway. In *Proceedings of the 7th Annual International Computing and Combinatorics Conference*, pages 237–246. LNCS, vol. 2108. Springer-Verlag, Berlin, 2001. Also to appear in *Theoretical Computer Science*.
2. J. H. Conway and N. J. A. Sloane. *Sphere Packings, Lattices and Groups*, 2nd edition. Springer-Verlag, New York, 1993.
3. H. Eiselt and G. Laporte. Competitive spatial models. *European Journal of Operational Research*, 39:231–242, 1989.
4. H. Eiselt, G. Laporte, and J.-F. Thisse. Competitive location models: a framework and bibliography. *Transportation Science*, 27:44–54, 1993.

5. G. Fejes Tóth. Sur la représentation d'une population infinie par une nombre fini d'elements. *Acta Mathematica Academiae Scientiarum Hungaricae*, 10:299–304, 1959.
6. L. Fejes Tóth. *Lagerungen in der Ebene, auf der Kugel und im Raum*, 2nd edition. Springer-Verlag, Berlin, 1972.
7. S. Fekete and H. Meijer. The one-round Voronoi game replayed. Manuscript, 2002.
8. B. Mohar and C. Thomassen. *Graphs on Surfaces*. Johns Hopkins University Press, Baltimore, MD, 2001.
9. R. Tobin, T. Friesz, and T. Miller. Existence theory for spatially competitive network facility location models. *Annals of Operations Research*, 18:267–276, 1989.

*Received July 3, 2002, and in revised form April 15, 2003. Online publication November 5, 2003.*