

Simple Rectilinear Polygons are Perfect under Rectangular Vision



Esther Arkin
Joseph S. B. Mitchell
Christiane Schmidt




Michael Biro




Omrit Filtser
Matthew J. Katz



The Art Gallery Problem, Guards and Witnesses




Rectangular Vision



Perfectness




Outlook



The Art Gallery Problem, Guards and Witnesses


The Art Gallery Problem, Guards and Witnesses



The Art Gallery Problem, Guards and Witnesses

Given: Polygon P

How many guards do we need to monitor P?




The Art Gallery Problem, Guards and Witnesses


Given: Polygon P

How many guards do we need to monitor P ?

minimum number of guards necessary for P : $g(P)$



The Art Gallery Problem, Guards and Witnesses



Given: Polygon P

How many guards do we need to monitor P ?

minimum number of guards necessary for P : $g(P)$


In general:

NP-hard


- point guards with holes [O'Rourke & Supowit 1983]
- vertex guards without holes [Lee & Lin 1986]
- point guards without holes [Aggarwal 1986]

APX-hard [Eidenbenz, Stamm & Widmayer 2001]

The Art Gallery Problem, Guards and Witnesses




The Art Gallery Problem, Guards and Witnesses



visibility-independent set of
witness points:

a finite set of points $W \subset P$,
whose visibility polygons are
pairwise disjoint.


The Art Gallery Problem, Guards and Witnesses



visibility-independent set of
witness points:

a finite set of points $W \subset P$,
whose visibility polygons are
pairwise disjoint.

The Art Gallery Problem, Guards and Witnesses




visibility-independent set of
witness points:

a finite set of points $W \subset P$,
whose visibility polygons are
pairwise disjoint.

$w(P)$: independence witness
number of P

The Art Gallery Problem, Guards and Witnesses




visibility-independent set of
witness points:

a finite set of points $W \subset P$,
whose visibility polygons are
pairwise disjoint.

$w(P)$: independence witness
number of P

Here: $w(P)=2$

The Art Gallery Problem, Guards and Witnesses




visibility-independent set of
witness points:

a finite set of points $W \subset P$,
whose visibility polygons are
pairwise disjoint.

$w(P)$: independence witness
number of P


Here: $w(P)=2$

polygon is **perfect** if $w(P)=g(P)$!




Rectangular Vision

Rectangular vision




Rectangular vision

- ordinary visibility:




Rectangular vision

- ordinary visibility:
 $p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .



Rectangular vision

- ordinary visibility:
 $p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .




Rectangular vision


- ordinary visibility:

$p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .

- **rectangular visibility:**



Rectangular vision




- ordinary visibility:

$p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .

- **rectangular visibility:**

$p, q \in P$ see each other if the rectangle spanned by those two points, $r(p, q)$, is fully contained in P .

Rectangular vision




- ordinary visibility:

$p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .

- **rectangular visibility:**

$p, q \in P$ see each other if the rectangle spanned by those two points, $r(p, q)$, is fully contained in P .

Rectangular vision




- ordinary visibility:

$p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .

- **rectangular visibility:**

$p, q \in P$ see each other if the rectangle spanned by those two points, $r(p, q)$, is fully contained in P .

Rectangular vision




- ordinary visibility:

$p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .

- **rectangular visibility:**

$p, q \in P$ see each other if the rectangle spanned by those two points, $r(p, q)$, is fully contained in P .

Rectangular vision



- ordinary visibility:


$p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .

- **rectangular visibility:**

$p, q \in P$ see each other if the rectangle spanned by those two points, $r(p, q)$, is fully contained in P .

Here: simple rectilinear polygons.

Rectangular vision



- ordinary visibility:

$p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .


- **rectangular visibility:**

$p, q \in P$ see each other if the rectangle spanned by those two points, $r(p, q)$, is fully contained in P .

Here: simple rectilinear polygons.

Notation:

Rectangular vision



- ordinary visibility:

$p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .

- **rectangular visibility:**


$p, q \in P$ see each other if the rectangle spanned by those two points, $r(p, q)$, is fully contained in P .

Here: simple rectilinear polygons.

Notation:

- visibility polygon of a point p in P : $VP(p)$

Rectangular vision



- ordinary visibility:

$p, q \in P$ see each other if the line segment connecting p and q is fully contained in P .

- **rectangular visibility:**


$p, q \in P$ see each other if the rectangle spanned by those two points, $r(p, q)$, is fully contained in P .

Here: simple rectilinear polygons.

Notation:


- visibility polygon of a point p in P : $VP(p)$
- sometimes: $VP_P(p)$

Rectangular vision



Rectangular vision

Known: for simple rectilinear polygon $g(P)$
can be computed in polynomial time - $O(n^{17})$:
[Worman&Keil, 2007]




Rectangular vision

Known: for simple rectilinear polygon $g(P)$
can be computed in polynomial time - $O(n^{17})$:

[Worman&Keil, 2007]

- extend all edges of reflex vertices of P




Rectangular vision

Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$:

[Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:




Rectangular vision

Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$:

[Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)




Rectangular vision

Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$:

[Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)
 - two nodes u and v are adjacent iff there is a star that contains both u and v




Rectangular vision

Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$:

[Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)
 - two nodes u and v are adjacent iff there is a star that contains both u and v




Rectangular vision

Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$:

[Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)
 - two nodes u and v are adjacent iff there is a star that contains both u and v




Rectangular vision

Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$:

[Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)
 - two nodes u and v are adjacent iff there is a star that contains both u and v




Rectangular vision


Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$:

[Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)
 - two nodes u and v are adjacent iff there is a star that contains both u and v




Rectangular vision



Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$: [Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)
 - two nodes u and v are adjacent iff there is a star that contains both u and v
- Theorem: Given a simple orthogonal polygon P and its $RVG(P)$, there is a star containing a set of regions U iff the members of U form clique in $RVG(P)$.


Rectangular vision



Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$: [Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)
 - two nodes u and v are adjacent iff there is a star that contains both u and v
- Theorem: Given a simple orthogonal polygon P and its $RVG(P)$, there is a star containing a set of regions U iff the members of U form clique in $RVG(P)$.
- Worman and Keil show that the class of region visibility graphs over all simple orthogonal polygons $\mathcal{RVG}$ are perfect graphs:


Rectangular vision



Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$: [Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)
 - two nodes u and v are adjacent iff there is a star that contains both u and v
- Theorem: Given a simple orthogonal polygon P and its $RVG(P)$, there is a star containing a set of regions U iff the members of U form clique in $RVG(P)$.
- Worman and Keil show that the class of region visibility graphs over all simple orthogonal polygons $\mathcal{RVG}$ are perfect graphs:
 - this allows for the computation of a minimum clique cover in polynomial time

Rectangular vision



Known: for simple rectilinear polygon $g(P)$ can be computed in polynomial time - $O(n^{17})$:


[Worman&Keil, 2007]

- extend all edges of reflex vertices of P
- build a graph (region visibility graph, $RVG(P)$) on that arrangement:
 - node per cell (=basic region)
 - two nodes u and v are adjacent iff there is a star that contains both u and v
- Theorem: Given a simple orthogonal polygon P and its $RVG(P)$, there is a star containing a set of regions U iff the members of U form clique in $RVG(P)$.
- Worman and Keil show that the class of region visibility graphs over all simple orthogonal polygons $\mathcal{RVG}$ are perfect graphs:
 - this allows for the computation of a minimum clique cover in polynomial time
 - proof: graphs in $\mathcal{RVG}$ have no large holes and no large odd antiholes (≥ 5)

Rectangular vision


Rectangular vision

- Computing a max independent witness set in P is equivalent to computing a max independent set in $RVG(P)$




Rectangular vision

- Computing a max independent witness set in P is equivalent to computing a max independent set in $RVG(P)$
- ➔ as $RVG(P)$ is perfect that also shows that $w(P)=g(P)$




Rectangular vision



- Computing a max independent witness set in P is equivalent to computing a max independent set in $RVG(P)$
 - ➔ as $RVG(P)$ is perfect that also shows that $w(P)=g(P)$
- we give a direct geometric proof (maybe simpler)

Rectangular Vision



Lemma (VPs):

Given three points $w, p, q \in P$. Then:


$$VP(w) \cap VP(p) \neq \emptyset$$

$$\wedge VP(w) \cap VP(q) \neq \emptyset$$

$$\wedge VP(q) \cap VP(p) \neq \emptyset$$

$$\Rightarrow VP(w) \cap VP(p) \cap VP(q) \neq \emptyset.$$

Rectangular Vision



Lemma (VPs):

Given three points $w, p, q \in P$. Then:


$$VP(w) \cap VP(p) \neq \emptyset$$

$$\wedge VP(w) \cap VP(q) \neq \emptyset$$

$$\wedge VP(q) \cap VP(p) \neq \emptyset$$

$$\Rightarrow VP(w) \cap VP(p) \cap VP(q) \neq \emptyset.$$

Rectangular Vision



Lemma (VPs):

Given three points $w, p, q \in P$. Then:


$$VP(w) \cap VP(p) \neq \emptyset$$

$$\wedge VP(w) \cap VP(q) \neq \emptyset$$

$$\wedge VP(q) \cap VP(p) \neq \emptyset$$

$$\Rightarrow VP(w) \cap VP(p) \cap VP(q) \neq \emptyset.$$

Rectangular Vision



Lemma (VPs):

Given three points $w, p, q \in P$. Then:


$$VP(w) \cap VP(p) \neq \emptyset$$

$$\wedge VP(w) \cap VP(q) \neq \emptyset$$

$$\wedge VP(q) \cap VP(p) \neq \emptyset$$

$$\Rightarrow VP(w) \cap VP(p) \cap VP(q) \neq \emptyset.$$

Rectangular Vision



Lemma (VPs):

Given three points $w, p, q \in P$. Then:


$$VP(w) \cap VP(p) \neq \emptyset$$

$$\wedge VP(w) \cap VP(q) \neq \emptyset$$

$$\wedge VP(q) \cap VP(p) \neq \emptyset$$

$$\Rightarrow VP(w) \cap VP(p) \cap VP(q) \neq \emptyset.$$

Rectangular Vision



Lemma (VPs):


Given three points $w, p, q \in P$. Then:

$$VP(w) \cap VP(p) \neq \emptyset$$

$$\wedge VP(w) \cap VP(q) \neq \emptyset$$

$$\wedge VP(q) \cap VP(p) \neq \emptyset$$

$$\Rightarrow VP(w) \cap VP(p) \cap VP(q) \neq \emptyset.$$




Not true for polygons with holes!




Theorem 1:

Simple rectilinear polygons are perfect under rectangular vision: $w(P)=g(P)$.



Perfectness





Perfectness





Perfectness





Perfectness



Perfectness



Perfectness




Perfectness

Proof by induction on the number, k , of witnesses:

Perfectness

Proof by induction on the number, k , of witnesses:


❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$



Perfectness

Proof by induction on the number, k , of witnesses:


❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$



Perfectness

Proof by induction on the number, k , of witnesses:

- ❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$
- ❖ **Proof:** Lemma on VPs + Theorem by Breen [Breen, 1996]




Proof by induction on the number, k , of witnesses:

- ❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$
- ❖ **Proof:** Lemma on VPs + Theorem by Breen [Breen, 1996]
- ❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)

Proof by induction on the number, k , of witnesses:

- ❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$
- ❖ **Proof:** Lemma on VPs + Theorem by Breen [Breen, 1996]
- ❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)
- ❖ **Induction Step $k \rightarrow k+1$:** Let P' be a polygon with $k+1$ witnesses.


Perfectness



Proof by induction on the number, k , of witnesses:

- ❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$
- ❖ **Proof:** Lemma on VPs + Theorem by Breen [Breen, 1996]
- ❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)
- ❖ **Induction Step $k \rightarrow k+1$:** Let P' be a polygon with $k+1$ witnesses.


Perfectness



Proof by induction on the number, k , of witnesses:

- ❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$
- ❖ **Proof:** Lemma on VPs + Theorem by Breen [Breen, 1996]
- ❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)
- ❖ **Induction Step $k \rightarrow k+1$:** Let P' be a polygon with $k+1$ witnesses.
- ❖ **Idea:** cover P' with 2 polygons:


Perfectness



Proof by induction on the number, k , of witnesses:

- ❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$
- ❖ **Proof:** Lemma on VPs + Theorem by Breen [Breen, 1996]
- ❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)
- ❖ **Induction Step $k \rightarrow k+1$:** Let P' be a polygon with $k+1$ witnesses.
- ❖ **Idea:** cover P' with 2 polygons:
 - ⊙ P_1 : 1-guardable

Perfectness



Proof by induction on the number, k , of witnesses:

❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$

❖ **Proof:** Lemma on VPs + Theorem by Breen [Breen, 1996]


❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)

❖ **Induction Step $k \rightarrow k+1$:** Let P' be a polygon with $k+1$ witnesses.

❖ **Idea:** cover P' with 2 polygons:

⊙ P_1 1-guardable

Perfectness



Proof by induction on the number, k , of witnesses:

❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$

❖ Proof: Lemma on VPs + Theorem by Breen [Breen, 1996]

❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)


❖ **Induction Step $k \rightarrow k+1$:** Let P' be a polygon with $k+1$ witnesses.

❖ Idea: cover P' with 2 polygons:

⊙ P_1 1-guardable

⊙ P_2 : a possibly disconnected polygon that has at most k witnesses w.r.t. visibility in P' (I.H.: is k -guardable)


Perfectness



Proof by induction on the number, k , of witnesses:

- ❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$
- ❖ **Proof:** Lemma on VPs + Theorem by Breen [Breen, 1996]
- ❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)
- ❖ **Induction Step $k \rightarrow k+1$:** Let P' be a polygon with $k+1$ witnesses.
- ❖ **Idea:** cover P' with 2 polygons:
 - ⊙ P_1 1-guardable
 - ⊙ P_2 : a possibly disconnected polygon that has at most k witnesses w.r.t. visibility in P' (I.H.: is k -guardable)

Perfectness



Proof by induction on the number, k , of witnesses:

❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$

❖ Proof: Lemma on VPs + Theorem by Breen [Breen, 1996]

❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)

❖ **Induction Step $k \rightarrow k+1$:** Let P' be a polygon with $k+1$ witnesses.


❖ Idea: cover P' with 2 polygons:

⊙ P_1 1-guardable

⊙ P_2 : a possibly disconnected polygon that has at most k witnesses w.r.t. visibility in P' (I.H.: is k -guardable)

➡ P' is $(k+1)$ -guardable

Perfectness



Proof by induction on the number, k , of witnesses:

❖ **Base Case:** $w(P)=1 \Rightarrow g(P)=1$

❖ Proof: Lemma on VPs + Theorem by Breen [Breen, 1996]

❖ **Induction Hypothesis:** $w(P)=k \Rightarrow g(P)=k$
(If P has k witnesses, P can be monitored using k guards.)

❖ **Induction Step $k \rightarrow k+1$:** Let P' be a polygon with $k+1$ witnesses.

❖ Idea: cover P' with 2 polygons:

⊙ P_1 1-guardable

⊙ P_2 : a possibly disconnected polygon that has at most k witnesses w.r.t. visibility in P' (I.H.: is k -guardable)

➔ P' is $(k+1)$ -guardable

How to construct these polygons?

We start with a specific witness:

We start with a specific witness:

- ❖ Claim: There exists an **ear witness**, i.e., a witness w such that all other witnesses $w_1, \dots, w_k$ are in the same connected component of $P' \setminus VP(w)$.

We start with a specific witness:

- ❖ Claim: There exists an **ear witness**, i.e., a witness w such that all other witnesses $w_1, \dots, w_k$ are in the same connected component of $P' \setminus VP(w)$.

Let this connected component be CC_k .

We start with a specific witness:

❖ Claim: There exists an **ear witness**, i.e., a witness w such that all other witnesses $w_1, \dots, w_k$ are in the same connected component of $P' \setminus VP(w)$.

Let this connected component be CC_k .

witness set: $W = \{w, w_1, \dots, w_k\}$

Perfectness


We start with a specific witness:


- ❖ Claim: There exists an **ear witness**, i.e., a witness w such that all other witnesses $w_1, \dots, w_k$ are in the same connected component of $P' \setminus VP(w)$.

Let this connected component be CC_k .

witness set: $W = \{w, w_1, \dots, w_k\}$

- ❖ Proof:





We start with a specific witness:

- ❖ Claim: There exists an **ear witness**, i.e., a witness w such that all other witnesses $w_1, \dots, w_k$ are in the same connected component of $P' \setminus VP(w)$.

Let this connected component be CC_k .

witness set: $W = \{w, w_1, \dots, w_k\}$

- ❖ Proof:

Consider the red/white decomposition: color visibility polygons of points in W red, the remaining subpolygons white.

Perfectness

We start with a specific witness:

- ❖ Claim: There exists an **ear witness**, i.e., a witness w such that all other witnesses $w_1, \dots, w_k$ are in the same connected component of $P' \setminus VP(w)$.


Let this connected component be CC_k .

witness set: $W = \{w, w_1, \dots, w_k\}$

- ❖ Proof:

Consider the red/white decomposition: color visibility polygons of points in W red, the remaining subpolygons white.

Its dual is a tree, and naturally 2-colored.




Perfectness

We start with a specific witness:

- ❖ Claim: There exists an **ear witness**, i.e., a witness w such that all other witnesses $w_1, \dots, w_k$ are in the same connected component of $P' \setminus VP(w)$.

Let this connected component be CC_k .

witness set: $W = \{w, w_1, \dots, w_k\}$




- ❖ Proof:

Consider the red/white decomposition: color visibility polygons of points in W red, the remaining subpolygons white.

Its dual is a tree, and naturally 2-colored.

Define a root arbitrarily,

Perfectness



We start with a specific witness:

- ❖ Claim: There exists an **ear witness**, i.e., a witness w such that all other witnesses $w_1, \dots, w_k$ are in the same connected component of $P' \setminus VP(w)$.

Let this connected component be CC_k .

witness set: $W = \{w, w_1, \dots, w_k\}$

- ❖ Proof:


Consider the red/white decomposition: color visibility polygons of points in W red, the remaining subpolygons white.

Its dual is a tree, and naturally 2-colored.

Define a root arbitrarily,

and take the lowest red vertex (red vertex who has, if any, white descendants)

Perfectness



We start with a specific witness:

- ❖ Claim: There exists an **ear witness**, i.e., a witness w such that all other witnesses $w_1, \dots, w_k$ are in the same connected component of $P' \setminus VP(w)$.

Let this connected component be CC_k .

witness set: $W = \{w, w_1, \dots, w_k\}$

- ❖ Proof:

Consider the red/white decomposition: color visibility polygons of points in W red, the remaining subpolygons white.

Its dual is a tree, and naturally 2-colored.

Define a root arbitrarily,


and take the lowest red vertex (red vertex who has, if any, white descendants)

witness corresponding to that vertex is an ear witness

- Idea: place a guard g in $VP(w)$, define $VP(g)$ as $P1$


- Idea: place a guard g in $VP(w)$, define $VP(g)$ as P_1
- AND: make sure that the remaining polygon does not allow more than k witnesses!

Perfectness




- Idea: place a guard g in $VP(w)$, define $VP(g)$ as P_1
- AND: make sure that the remaining polygon does not allow more than k witnesses!

Perfectness




- Idea: place a guard g in $VP(w)$, define $VP(g)$ as P_1
- AND: make sure that the remaining polygon does not allow more than k witnesses!

Perfectness




- Idea: place a guard g in $VP(w)$, define $VP(g)$ as P_1
- AND: make sure that the remaining polygon does not allow more than k witnesses!

Perfectness



- Idea: place a guard g in $VP(w)$, define $VP(g)$ as P_1
- AND: make sure that the remaining polygon does not allow more than k witnesses!


Perfectness



- Idea: place a guard g in $VP(w)$, define $VP(g)$ as P_1
- AND: make sure that the remaining polygon does not allow more than k witnesses!


Perfectness

$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$



Perfectness


$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$



Perfectness

$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$


- set of „private“ neighbors of w



Perfectness

$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$


- set of „private“ neighbors of w



Perfectness

$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$

- set of „private“ neighbors of w




Perfectness

$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$

- set of „private“ neighbors of w

$$Q = \cap_{u \in U} VP(u), Q \neq \emptyset$$




Perfectness

$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$

- set of „private“ neighbors of w

$$Q = \cap_{u \in U} VP(u), Q \neq \emptyset$$

$$(U = \emptyset : Q = VP(w))$$



Perfectness


$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$

- set of „private“ neighbors of w

$$Q = \cap_{u \in U} VP(u), Q \neq \emptyset$$

$$(U = \emptyset : Q = VP(w))$$

placing g in Q is still not enough!



Perfectness


$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$

- set of „private“ neighbors of w

$$Q = \cap_{u \in U} VP(u), Q \neq \emptyset$$

$$(U = \emptyset : Q = VP(w))$$

placing g in Q is still not enough!



Perfectness


$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$

- set of „private“ neighbors of w

$$Q = \cap_{u \in U} VP(u), Q \neq \emptyset$$

$$(U = \emptyset : Q = VP(w))$$

placing g in Q is still not enough!



Perfectness


$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$

- set of „private“ neighbors of w

$$Q = \cap_{u \in U} VP(u), Q \neq \emptyset$$

$$(U = \emptyset : Q = VP(w))$$

placing g in Q is still not enough!



Perfectness

$$U = \{u \notin VP(w) : VP(u) \cap VP(w) \neq \emptyset; \\ VP(u) \cap VP(w_i) = \emptyset \forall i\}$$


- set of „private“ neighbors of w

$$Q = \cap_{u \in U} VP(u), Q \neq \emptyset$$

$$(U = \emptyset : Q = VP(w))$$

placing g in Q is still not enough!

Intuition: we need to match witness VPs from both CC_k and $P \setminus \{CC_k \cup VP(w)\}$



Perfectness

Helpful Lemma (IntersectionGraph):

Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.

Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.

Let G be the intersection graph of the visibility polygons of witnesses from W and W' .


Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.

Let G be the intersection graph of the visibility polygons of witnesses from W and W' .

Then any connected component of G contains an equal number of witnesses from W and W' .

Perfectness




Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.

Let G be the intersection graph of the visibility polygons of witnesses from W and W' .

Then any connected component of G contains an equal number of witnesses from W and W'

Perfectness




Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.

Let G be the intersection graph of the visibility polygons of witnesses from W and W' .

Then any connected component of G contains an equal number of witnesses from W and W'

Perfectness




Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.

Let G be the intersection graph of the visibility polygons of witnesses from W and W' .

Then any connected component of G contains an equal number of witnesses from W and W' .

Perfectness



Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.

Let G be the intersection graph of the visibility polygons of witnesses from W and W' .

Then any connected component of G contains an equal number of witnesses from W and W' .


Perfectness

Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.

Let G be the intersection graph of the visibility polygons of witnesses from W and W' .

Then any connected component of G contains an equal number of witnesses from W and W' .



Take 3 witnesses from W ,


Perfectness

Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.

Let G be the intersection graph of the visibility polygons of witnesses from W and W' .

Then any connected component of G contains an equal number of witnesses from W and W'



Take 3 witnesses from W ,
and the remaining $k-2$ from W'


Perfectness

Helpful Lemma (IntersectionGraph):

Let W and W' be two maximum cardinality witness sets for P , $w(P)=k$.


Let G be the intersection graph of the visibility polygons of witnesses from W and W' .

Then any connected component of G contains an equal number of witnesses from W and W' .




Take 3 witnesses from W ,
and the remaining $k-2$ from W'
➡ $>k$ witnesses

Perfectness




Perfectness



CC_k


Perfectness



$$P' \setminus \{CC_k \cup VP(w)\}$$

$$CC_k$$


Perfectness



$$P' \setminus \{CC_k \cup VP(w)\}$$


$$CC_k$$

Perfectness




$$P' \setminus \{CC_k \cup VP(w)\} \quad CC_k$$

Perfectness




Perfectness



Perfectness


For any IS X of visibility polygons
in $I_1 \cup I'_2$:



Perfectness

For any IS X of visibility polygons
in $I_1 \cup I'_2$:

$$|\{v \in W : VP(v) \cap VP(x), x \in X\}| \geq |X|$$




Perfectness

For any IS X of visibility polygons
in $I_1 \cup I'_2$:

$$|\{v \in W : VP(v) \cap VP(x), x \in X\}| \geq |X|$$

If X is a max IS of visibility
polygons in $I_1 \cup I'_2$ of size t




Perfectness

For any IS X of visibility polygons in $I_1 \cup I'_2$:

$$|\{v \in W : VP(v) \cap VP(x), x \in X\}| \geq |X|$$

If X is a max IS of visibility polygons in $I_1 \cup I'_2$ of size t

➡ exists a matching of size t in the intersection graph of VPs from points in W and points in X



Perfectness


For any IS X of visibility polygons in $I_1 \cup I'_2$:

$$|\{v \in W : VP(v) \cap VP(x), x \in X\}| \geq |X|$$

If X is a max IS of visibility polygons in $I_1 \cup I'_2$ of size t

→ exists a matching of size t in the intersection graph of VPs from points in W and points in X

Let the stabbing number of visibility polygons from $I_1 \cup I'_2$ be s .



Perfectness

For any IS X of visibility polygons in $I_1 \cup I'_2$:


$$|\{v \in W : VP(v) \cap VP(x), x \in X\}| \geq |X|$$

If X is a max IS of visibility polygons in $I_1 \cup I'_2$ of size t

→ exists a matching of size t in the intersection graph of VPs from points in W and points in X

Let the stabbing number of visibility polygons from $I_1 \cup I'_2$ be s .

$s=t$:



Perfectness

For any IS X of visibility polygons in $I_1 \cup I'_2$:

$$|\{v \in W : VP(v) \cap VP(x), x \in X\}| \geq |X|$$


If X is a max IS of visibility polygons in $I_1 \cup I'_2$ of size t

→ exists a matching of size t in the intersection graph of VPs from points in W and points in X

Let the stabbing number of visibility polygons from $I_1 \cup I'_2$ be s .

$s=t$:

- s_1 polygons that do not interfere



Perfectness

For any IS X of visibility polygons in $I_1 \cup I'_2$:

$$|\{v \in W : VP(v) \cap VP(x), x \in X\}| \geq |X|$$


If X is a max IS of visibility polygons in $I_1 \cup I'_2$ of size t

➔ exists a matching of size t in the intersection graph of VPs from points in W and points in X

Let the stabbing number of visibility polygons from $I_1 \cup I'_2$ be s .

$s=t$:

- s_1 polygons that do not interfere
- $s_2 = s - s_1$ remaining:



Perfectness

For any IS X of visibility polygons in $I_1 \cup I'_2$:

$$|\{v \in W : VP(v) \cap VP(x), x \in X\}| \geq |X|$$


If X is a max IS of visibility polygons in $I_1 \cup I'_2$ of size t

➔ exists a matching of size t in the intersection graph of VPs from points in W and points in X

Let the stabbing number of visibility polygons from $I_1 \cup I'_2$ be s .

$s=t$:

- s_1 polygons that do not interfere
- $s_2 = s - s_1$ remaining:
 - stabbing number = minimum edge cover in the bipartite intersection graph



Perfectness

For any IS X of visibility polygons in $I_1 \cup I'_2$:

$$|\{v \in W : VP(v) \cap VP(x), x \in X\}| \geq |X|$$


If X is a max IS of visibility polygons in $I_1 \cup I'_2$ of size t

➡ exists a matching of size t in the intersection graph of VPs from points in W and points in X

Let the stabbing number of visibility polygons from $I_1 \cup I'_2$ be s .

$s=t$:

- s_1 polygons that do not interfere
 - $s_2 = s - s_1$ remaining:
 - stabbing number = minimum edge cover in the bipartite intersection graph
- ➡ max IS of size s




Perfectness

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.


Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.



Perfectness


Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.



Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$




Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.




Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$




Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$




Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$



Perfectness


Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.



Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$


❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W' with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be in the same connected component of the intersection graph (VP 's from W and W') as w



Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$


❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be in the same connected component of the intersection graph (VP 's from W and W') as w



Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$


❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be in the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r



Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

❖ $s=r-1$:

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be int the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be in the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r

❖ $s=r-1$:

- $k+1-s$ witnesses of W' are in a different connected component

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be in the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r

❖ $s=r-1$:

- $k+1-s$ witnesses of W' are in a different connected component

➡ $w, w_1, \dots, w_{r-1}, w'_{s+1}, \dots, w'_{k+1}$ form independent witness set of size $k+2$

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be in the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r

❖ $s=r-1$:

- $k+1-s$ witnesses of W' are in a different connected component

➡ $w, w_1, \dots, w_{r-1}, w'_{s+1}, \dots, w'_{k+1}$ form independent witness set of size $k+2$

❖ $r < s$:

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be in the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r

❖ $s=r-1$:

- $k+1-s$ witnesses of W' are in a different connected component

➡ $w, w_1, \dots, w_{r-1}, w'_{s+1}, \dots, w'_{k+1}$ form independent witness set of size $k+2$

❖ $r < s$:

➡ $w'_1, \dots, w'_s, w_r, \dots, w_k$ are an independent witness set $> k+1$

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be int the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r

❖ $s=r-1$:

- $k+1-s$ witnesses of W' are in a different connected component

➡ $w, w_1, \dots, w_{r-1}, w'_{s+1}, \dots, w'_{k+1}$ form independent witness set of size $k+2$

❖ $r < s$:

➡ $w'_1, \dots, w'_s, w_r, \dots, w_k$ are an independent witness set $> k+1$

❖ $r=s$:

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be int the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r

❖ $s=r-1$:

- $k+1-s$ witnesses of W' are in a different connected component

➡ $w, w_1, \dots, w_{r-1}, w'_{s+1}, \dots, w'_{k+1}$ form independent witness set of size $k+2$

❖ $r < s$:

➡ $w'_1, \dots, w'_s, w_r, \dots, w_k$ are an independent witness set $> k+1$

❖ $r=s$:

- we use as many witnesses in the connected component of w as are maximally stabbed

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be int the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r

❖ $s=r-1$:

- $k+1-s$ witnesses of W' are in a different connected component

➔ $w, w_1, \dots, w_{r-1}, w'_{s+1}, \dots, w'_{k+1}$ form independent witness set of size $k+2$

❖ $r < s$:

➔ $w'_1, \dots, w'_s, w_r, \dots, w_k$ are an independent witness set $> k+1$

❖ $r=s$:

- we use as many witnesses in the connected component of w as are maximally stabbed

➔ one of $w'_1, \dots, w'_s$ is in P_1

Perfectness

Place guard g in Q : monitoring $VP(w)$ and all points assigned via stabbing.

$$P_1 = VP(g)$$

❖ P_1 is 1-guardable.

$$P_2 = \bigcup_{i=1}^k VP(w_i) \cup \left(\bigcup_{i=1}^k VP(VP(w_i)) \setminus P_1 \right)$$

❖ $w(P_2) \leq k$.

❖ Proof by contradiction:

- Assume there is a witness set $W' = \{w'_1, \dots, w'_{k+1}\}$ of size $k+1$ in P_2
- Consider all w'_i in W with $VP_{P'}(w'_i) \cap VP_{P'}(w) \neq \emptyset$
- w.l.o.g. let $w'_1, \dots, w'_s$ be in the same connected component of the intersection graph (VP 's from W and W') as w
- number of witnesses from W in this connected component: r

❖ $s=r-1$:

- $k+1-s$ witnesses of W' are in a different connected component

➡ $w, w_1, \dots, w_{r-1}, w'_{s+1}, \dots, w'_{k+1}$ form independent witness set of size $k+2$

❖ $r < s$:

➡ $w'_1, \dots, w'_s, w_r, \dots, w_k$ are an independent witness set $> k+1$

❖ $r=s$:


- we use as many witnesses in the connected component of w as are maximally stabbed

➡ one of $w'_1, \dots, w'_s$ is in P_1

Finally: $P' = P_1 \cup P_2$




Outlook




Outlook

- computation of $w(P)$ in substantiell lower running time ($<O(n^{17})$) via Dynamic Programming





Outlook



- computation of $w(P)$ in substantiell lower running time ($<O(n^{17})$) via Dynamic Programming
- ➔ faster algorithm to compute $g(P)$


Outlook



- computation of $w(P)$ in substantially lower running time ($<O(n^{17})$) via Dynamic Programming
- ➔ faster algorithm to compute $g(P)$
- ➔ as the above is constructive: also computation of the location of guards possible?



- computation of $w(P)$ in substantiell lower running time ($<O(n^{17})$) via Dynamic Programming
- ➡ faster algorithm to compute $g(P)$
- ➡ as the above is constructive: also computation of the location of guards possible?
- other types of vision?

Outlook




- computation of $w(P)$ in substantiell lower running time ($<O(n^{17})$) via Dynamic Programming
- ➔ faster algorithm to compute $g(P)$
- ➔ as the above is constructive: also computation of the location of guards possible?
- other types of vision?
 - „staircase“ visibility

Outlook




- computation of $w(P)$ in substantially lower running time ($<O(n^{17})$) via Dynamic Programming
- ➔ faster algorithm to compute $g(P)$
- ➔ as the above is constructive: also computation of the location of guards possible?
- other types of vision?
 - „staircase“ visibility
 - ordinary visibility

Thanks.




The image features a light blue background with dark blue horizontal bands at the top and bottom. A central composition of overlapping rectangles is outlined in dark grey. The rectangles include a large light green one in the center, a blue one at the top left, a yellow-green one at the top center, a purple one at the top right, and a teal one at the top far right. On the left side, there are teal, yellow-green, and blue rectangles. A grey rectangle is positioned below the purple one. Four circles are placed within the composition: a blue circle in the bottom-left green rectangle, a yellow circle in the center green rectangle, a pink circle in the purple rectangle, and a green circle in the teal rectangle. The word "Thanks." is written in white, bold, sans-serif font inside a white rectangular box with a dark border, centered over the main green rectangle.


Thanks.




Computation of $w(P)$




The Art Gallery Problem, Guards and Witnesses



Rectangular Vision



Perfectness




Computation of $w(P)$



Outlook

Thanks.




Thanks.


Theorem by Breen [1996]:

Let $\mathcal{P}$ be a family of simple polygons in the plane.

- If every three members of $\mathcal{P}$ have a simply connected union
 - and every two have a nonempty intersection
- ➡ $\cap\{P : P \in \mathcal{P}\} \neq \emptyset$
- for a family of orthogonally convex polygons the intersection is again an orthogonally convex polygon.




Known:



Known:


Simple polygon with n vertices: $\lfloor n/3 \rfloor$ are sometimes necessary and always sufficient. [Chvátal '75]



Known:

Simple polygon with n vertices: $\lfloor n/3 \rfloor$ are sometimes necessary and always sufficient. [Chvátal '75]

Polygon with n vertices and h holes: $\lfloor (n + h)/3 \rfloor$ are sometimes necessary and always sufficient. [Bjorling-Sachs, Souvaine '95]




Known:

Simple polygon with n vertices: $\lfloor n/3 \rfloor$ are sometimes necessary and always sufficient. [Chvátal '75]

Polygon with n vertices and h holes:
 $\lfloor (n + h)/3 \rfloor$ are sometimes necessary and always sufficient. [Bjorling-Sachs, Souvaine '95]

NP-hard




Known:

Simple polygon with n vertices: $\lfloor n/3 \rfloor$ are sometimes necessary and always sufficient. [Chvátal '75]

Polygon with n vertices and h holes:
 $\lfloor (n + h)/3 \rfloor$ are sometimes necessary and always sufficient. [Bjorling-Sachs, Souvaine '95]

NP-hard

- point guards with holes [O'Rourke & Supowit 1983]




Known:

Simple polygon with n vertices: $\lfloor n/3 \rfloor$ are sometimes necessary and always sufficient. [Chvátal '75]

Polygon with n vertices and h holes:
 $\lfloor (n + h)/3 \rfloor$ are sometimes necessary and always sufficient. [Bjorling-Sachs, Souvaine '95]

NP-hard

- point guards with holes [O'Rourke & Supowit 1983]
- vertex guards without holes [Lee & Lin 1986]




Known:

Simple polygon with n vertices: $\lfloor n/3 \rfloor$ are sometimes necessary and always sufficient. [Chvátal '75]

Polygon with n vertices and h holes:
 $\lfloor (n + h)/3 \rfloor$ are sometimes necessary and always sufficient. [Bjorling-Sachs, Souvaine '95]

NP-hard

- point guards with holes [O'Rourke & Supowit 1983]
- vertex guards without holes [Lee & Lin 1986]
- point guards without holes [Aggarwal 1986]



Known:

Simple polygon with n vertices: $\lfloor n/3 \rfloor$ are sometimes necessary and always sufficient. [Chvátal '75]

Polygon with n vertices and h holes: $\lfloor (n + h)/3 \rfloor$ are sometimes necessary and always sufficient. [Bjorling-Sachs, Souvaine '95]

NP-hard

- point guards with holes [O'Rourke & Supowit 1983]
- vertex guards without holes [Lee & Lin 1986]
- point guards without holes [Aggarwal 1986]

APX-hard [Eidenbenz, Stamm & Widmayer 2001]

